

UNCLASSIFIED

AD 410524

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION


CAMERON STATION, ALEXANDRIA, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

3



393500

14-63-4-3

410524

See note b.

1 NA

2 NA

3 NA

4 See note b.

5 4 April 63,

6 6 p.

7 NA

8 NA

9 NA

10 NA

11 NA

12 U


13 NA

Q12

11/1/63

AD No. _____


DDC FILE COPY



410524

REC'D 12695
Report Number

2000



JUL 30 1963
JISIA A

United States Atomic Energy Commission
Division of Technical Information

APR 4 1954

Constant in HD

J. I. Musher

Department of Chemistry, Harvard University
Cambridge 38, Massachusetts

NP-12695

MASTER

(Received

The coupling of nuclear spins by second-order perturbation mechanisms was first proposed and discussed by Ramsey and Purcell¹ in explanation of experimental results obtained by Hahn and Maxwell.² Invoking an arbitrary "average energy" denominator to compare magnitudes of the several terms for the HD molecule, Ramsey³ showed that the dominant contribution to the indirect spin-spin coupling arises from the second-order perturbation theory energy of the Fermi-contact Hamiltonian,⁴ $\mathcal{H}_F = (16\pi\beta\hbar/3) \sum_{k,N} \gamma_N \delta(r_{kN}) S_k \cdot I_N$, where $r_{kN} = r_k - r_N$ and S_k and I_N are the electron and nuclear spins respectively. Stephen⁵ and others⁶ have since performed variational calculations for this same energy.

A new approximate procedure⁷ for performing perturbation calculations on simple unperturbed molecular wave-functions introducing no arbitrary parameters, enables the relatively facile calculation of some molecular properties.⁸ This procedure is used here to calculate approximately the Fermi-contact contribution,

* This research was supported in part by a fellowship from the Division of General Medical Sciences, Public Health Service and by support extended Harvard University by the Office of Naval Research.

J_F to the indirect scalar coupling between the nuclear spins in HD. The scalar coupling constant is written as J , where the coupling energy is $E = \hbar J I_H \cdot I_D + \hbar I_H \cdot g I_D$ and g is a traceless tensor whose effect vanishes under the usual experimental conditions. Consider the ground-state described by the Coulson-MO wave-function,⁹ $|0(r_1, r_2), 0(\sigma_1, \sigma_2)\rangle = 2^{-\frac{1}{2}} N^{-2} |(H+D)(1)(H+D)(2)[\alpha(1)\beta(2)-\alpha(2)\beta(1)]\rangle$ assuming the space- and spin-coordinates be separable for the excited states as well. $|H(1)\rangle$ and $|D(1)\rangle$ are one-electron $1s$ AO's centered on the proton and the deuteron respectively with, e.g. $|H(1)\rangle = (\alpha^3/\pi)^{\frac{1}{2}} \exp(-\alpha r_{1H})$, $\alpha = 1.197$, $N^2 = 2(1 + \langle A|B\rangle)$ and $R_{HD} = 1.4217$ a.u. J_F can be written, considering both the one-electron and two-electron contributions, as³

$$J_F = 2(2/3\hbar)(16\pi\beta\hbar/3)^2 \gamma_H \gamma_D \sum' (E_0 - E_n)^{-1} \{ \langle 0,0 | \delta(r_{1H}) S_1 | n,n \rangle \cdot \langle n,n | \delta(r_{1D}) S_1 | 0,0 \rangle + \langle 0,0 | \delta(r_{2H}) S_2 | n,n \rangle \cdot \langle n,n | \delta(r_{1D}) S_1 | 0,0 \rangle \} \quad (1)$$

The "covalent approximation" observes⁷ that if $\mathcal{H}_0 |0,0\rangle \equiv E_0 |0,0\rangle$ (\equiv since $|0,0\rangle$ is never an exact eigenfunction of \mathcal{H}_0), then $\mathcal{H}_0 |A(1)(A+B)(2), 0\rangle \approx E_0 |A(1)(A+B)(2), 0\rangle$ and $\mathcal{H}_0 |B(1)(A+B)(2), 0\rangle \approx E_0 |B(1)(A+B)(2), 0\rangle$. Let the function $f_N[\delta(r_{1N})]$ for each electron i and nuclei N, N' , be defined by the inhomogeneous p.d.e.

$$[f_N \{ \delta(r_{1N'}) \} \mathcal{H}_0] |N(1)\rangle = \delta(r_{1N'}) |N(1)\rangle - \langle N(1) | \delta(r_{1N'}) | N(1) \rangle |N(1)\rangle \quad (2)$$

The spatial function for electron $j \neq 1$ and the spin functions have been suppressed, defining $f\{\delta(r_1)\}$ to be independent of r_j . For $N = N'$ the solution to Eq. (2) is¹⁰

$$f_N\{\delta(r_{1N})\} = (2m/4\pi\hbar^2)(-1/r_{1N} + 2 \propto \ln r_{1N} + 2\alpha r_{1N}) \quad (3)$$

We have not succeeded in solving for $f_N\{\delta(r_{1N'})\}$, $N \neq N'$, although these f 's will be small as the inhomogeneities themselves, e.g. $\delta(r_H) |D\rangle \sim \exp(-\alpha R_{HD}) = 0.19$, are small. Their contributions are neglected as argued below. Following Dalgarno¹¹ and Schwartz¹⁰ the commutators of the f 's are substituted into Eq. (1) neglecting the non-orthogonality of $|A\rangle$ and $|B\rangle$ with $|n\rangle$ which gives a small error,⁷ and the sum over $|n\rangle$ is taken, observing that $\langle 0 | S_1^2 | 0 \rangle = -\langle 0 | S_1 \cdot S_2 | 0 \rangle = \frac{3}{4}$ and $\langle 0 | S_1 | 0 \rangle = 0$. J_F then becomes¹²

$$J_F = \hbar^{-1}(16\pi\hbar^2/3)^2 \gamma_H \gamma_D N^{-4} \left\{ N^2 \left[\langle (H+D)(1) | \delta(r_{1H}) f_D \{ \delta(r_{1D}) \} \right. \right. \\ \left. \left. |D(1)\rangle \right] + \langle H(1) | f_H \{ \delta(r_{1H}) \} \delta(r_{1D}) | H(1)\rangle + X_1 \right] - \\ \left[\langle (H+D)(1)(H+D)(2) | \delta(r_{1D}) f_D \{ \delta(r_{1D}) \} | D(1)(H+D)(2)\rangle \right. \\ \left. + \langle (H+D)(1)H(2) | f_H \{ \delta(r_{2H}) \} \delta(r_{1D}) | H(1)(H+D)(2)\rangle + X_2 \right] \} \quad (4)$$

with X_1 given by $X_1 = \Sigma' (E_0 - E_n)^{-1} \langle D(1)(H+D)(2) | \delta(r_{1H}) | n \rangle \langle n | \delta(r_{1D}) | H(1)(H+D)(2) \rangle$ and X_2 analogously. Such terms will clearly be small as both matrix elements involve the value at one nucleus of the AO centered on the other, being of the order of $\exp(-2\alpha R_{HD})$. X_1 and X_2 should each be $\exp(-\alpha R_{HD})$ smaller than the next largest one-electron and two-electron terms respectively,

which are themselves $\exp(-\alpha R)$ smaller than the respective dominant terms. The divergence of the Fermi self-coupling (J_F, HH or J_F, DD), when nuclear size corrections are neglected, is observed trivially.

When the necessary integrals are evaluated, the calculated contribution to J_F from the one-electron terms is 3.0 cps while that from the two-electron terms is -32.5 cps for a total of -29.5 cps. The error from both the method and the wave function (e.g., too much ionic contribution) is probably within 25% and will be discussed elsewhere along with results of other calculations. Experiment¹³ gives $J_{HD} = 43$ cps of indeterminate sign, and the calculated J_F indicates that indeed J_F is a principal contributor to J_{HD} . (The different orbital contributions³ will be discussed elsewhere.) The sign of J_{HD} could, in principle, be determined by ultra-high-resolution molecular beam techniques.

The negative calculated J_F is surprising since the approximation using an "average energy" denominator with the usual positive (and arbitrary) ΔE^* gives³ J_{HD} incorrectly as positive. This exhibits the well-known fallibility¹⁴ of the "average energy" procedure for sums whose numerators are not positive-definite.

The negative J_F can be argued physically. Consider the two-electron term, the energy of which is expressed as proportional to $(I_H S_1)(I_D S_2)$ whose dominant contribution has electron 1 on H and electron 2 on D. If H and D have their spins parallel, then by the Pauli principle, either $I_H S_1$ is positive and $I_D S_2$ negative, or vice versa, and the energy which is their product is negative. Since, for the scalar coupling, $E = \hbar^2 J_{HD} I_H I_D$, J_{HD}

is negative.¹⁵ The quantum mechanical argument would observe that most of the lowest coupled triplet states, rather than being "anti-bonding" states, e.g. $|\mathcal{A}(H-D)(1)(H+D)(2)\rangle$, are "bonding" states of higher AO's, e.g. $|\mathcal{A}(H+D)(1)(H'+D')(2)\rangle$ where H', D' are higher S_- states. For such $|n\rangle$ in the (dominant) two-electron sum, whenever the l.h. matrix element is positive, the r.h. one is negative. The product is therefore negative, which with a negative energy denominator and a negative spin matrix element, gives J_{HD} negative.

REFERENCES

1. N. F. Ramsey and E. M. Purcell, Phys. Rev. 85, 143 (1952).
2. E. L. Hahn and D. E. Maxwell, Phys. Rev. 84, 1246 (1951), and 88, 243 (1952). H. G. Gutowsky, D. McCall and C.P. Slichter, Phys. Rev. 84, 589 (1951).
3. N. F. Ramsey, Phys. Rev. 91, 303 (1953).
4. E. Fermi, Z. Physik 60, 320 (1930).
5. M. J. Stephen, Proc. Roy. Soc. A243, 274 (1957).
6. T. P. Das and R. Bersohn, Phys. Rev. 115, 897 (1959); D. E. O'Reilly, J. Chem. Phys. 36, 274 (1962).
7. J. I. Musher, submitted to Ann. Phys. (N.Y.)
8. J. I. Musher, Disc. Faraday Soc. (in press).
9. C. A. Coulson, Trans. Faraday Soc. 33, 1479 (1937).
10. C. Schwartz, Ann. Phys. (N.Y.) 2, 156, 170, 178 (1959).
11. A. Dalgarno and J. T. Lewis, Proc. Roy. Soc. A233, 70 (1955).
12. This is similar to Eq. (62) of ref. 6, but a different variational function is used for each center analogous to variationally determined screening in simple H_2 wave functions. Calculation shows the important term of Eq. (3) to be the one in r_{1N} . Therefore variational calculations using $r_H |H\rangle + r_D |D\rangle$ as a trial function would have met with better success than refs. 5 and 6.
13. B. Smaller, E. Vasaitis, E. C. Avery and D. A. Hutchinson, Phys. Rev. 88, 414 (1952); H. Y. Carr and E. M. Purcell, Phys. Rev. 88, 415 (1952).
14. A. D. McLachlan, J. Chem. Phys. 32, 1263 (1960).
15. This should not be looked at, as has been done in the past, from the point of view of additive energies in S_1 , as these are the first-order hf energies which equal to zero when both electrons are included.

END